

PRODUCT INFORMATION



PERFORMANCE MINERALS®

ZINPRO® 40

For use only in U.S.


ZINC METHIONINE COMPLEX

A nutritional feed ingredient for livestock and poultry. When used as a commercial feed ingredient it must be declared as zinc methionine complex. Association of American Feed Control Officials (AAFCO) No. 57.151. International Feed Number (IFN) 6-16-293. Zinc Methionine Complex. U.S. Patent No. 4,764,633 and 5,430,164. Canada Reg. No. 990581.

TYPICAL ANALYSIS

Zinc 4.0%

MOLECULAR STRUCTURE



PHYSICAL DESCRIPTION

A light brown granular powder with characteristic methionine-like odor. ZINPRO® 40 weighs approximately 32 lb/cu. ft. and is packaged in 25 kg (55.1 lb) multiwall bags.

FEEDING INSTRUCTIONS

SWINE*: Add 2.5 lb (1,134 grams) per ton of complete diet.

LAYING HENS, BROILERS AND TURKEYS: Add 2 lb (908 grams) per ton of complete diet.

DAIRY CATTLE: Feed 9.0 grams per head daily, or 2 lb per 100 head daily.

BEEF CATTLE: Feed 9.0 grams per head daily, or 2 lb per 100 head daily.

HORSES: Feed 9.0 grams per head daily.

SHEEP*: Feed 1.0 gram per head daily, or 1 lb per 454 head daily.

GOATS*: Feed 1.25 grams per head daily, or 1 lb per 360 head daily.

* Double recommended levels during periods of reduced feed intake, or high production stress.

TOXICITY AND SAFE HANDLING

When correctly used, there is no toxicity hazard in the use of ZINPRO 40. Store in a clean and dry environment. Follow label directions and rotate inventory to ensure fresh product.